arXiv:hep-ph/0406223v1 21 Jun 2004

Acta Phys. Hung. A 19/1 (2004) 000–000

**HEAVY ION PHYSICS**

# Unintegrated parton distributions and meson production in hadronic collisions

Antoni Szczurek[1,2] and Marta Czech[3]

[1] Institute of Nuclear Physics, PAN
PL-31-342 Cracow, Poland
[2] Rzeszów University
PL-35-959 Rzeszów, Poland
[3] M. Smoluchowski Institute of Physics
Jagiellonian University
PL-30-059 Cracow, Poland

*Received 1 January 2004*

**Abstract.** The inclusive distributions of gluons and pions for high-energy NN collisions are calculated. The results for several unintegrated gluon distributions (UGD's) from the literature are compared. We find huge differences in both rapidity and $p_t$ of gluons and $\pi$'s in NN collisions for different models of UGD's. The Karzeev-Levin UGD gives good description of momentum distribution of charged hadrons at midrapidities. We find, however, that the gluonic mechanism discussed does not describe the inclusive distributions of charged particles in the fragmentation region. Some of the missing mechanisms are calculated with the help of unintegrated parton distributions being solutions of the Kwieciński CCFM equations. We predict an asymmetry in $\pi^+$ and $\pi^-$ production which could be studied by the BRAHMS collaboration at RHIC.

*Keywords:* pion production, nucleon-nucleon collisions, unintegrated parton distributions, fragmentation functions, inclusive distributions
*PACS:* 12.38.Bx, 13.85.Hd, 13.85.Ni

## 1. Introduction

The recent results from RHIC (see e.g. [ 1]) have attracted renewed interest in better understanding the dynamics of particle production, not only in nuclear collisions. Quite different approaches have been used to describe the particle spectra from

1219-7580/04/ $ 20.00
© 2004 Akadémiai Kiadó, Budapest

the nuclear collisions [ 2]. The model in Ref.[ 3] with an educated guess for UGD describes surprisingly well the whole charged particle rapidity distribution by means of gluonic mechanisms only. Such a gluonic mechanism would lead to the identical production of positively and negatively charged hadrons. The recent results of the BRAHMS experiment concerning heavy ion collisions [ 4] put into question the successful description of Ref.[ 3]. In the light of this experiment, it becomes obvious that the large rapidity regions have more complicated flavour structure.

We discuss the relation between UGD's in hadrons and the inclusive momentum distribution of particles produced in hadronic collisions. The results obtained with different UGD's [ 3, 5, 7, 8, 9] are shown and compared.

In addition to the $gg \to g$ mechanism we include new mechanisms $q_f g \to q_f$ and $g q_f \to q_f$ and similar for antiquarks. We present first results based on unintegrated parton (gluon, quark, antiquark) distributions obtained by solving a set of coupled equations developed recently by Kwieciński and collaborators.

## 2. Inclusive gluon production

At sufficiently high energy the cross section for inclusive gluon production in $h_1 + h_2 \to g$ can be written in terms of the UGD's "in" both colliding hadrons [ 10]

$$\frac{d\sigma}{dy d^2p_t} = \frac{16N_c}{N_c^2-1}\frac{1}{p_t^2}\int \alpha_s(\Omega^2)\mathcal{F}_1(x_1,\kappa_1^2)\mathcal{F}_2(x_2,\kappa_2^2)\delta(\vec{\kappa}_1+\vec{\kappa}_2-\vec{p}_t)\; d^2\kappa_1 d^2\kappa_2 \; . \quad (1)$$

Above $\mathcal{F}_1$ and $\mathcal{F}_2$ are UGD's in hadron $h_1$ and $h_2$, respectively. The longitudinal momentum fractions are fixed by kinematics: $x_{1/2} = \frac{p_t}{\sqrt{s}} \cdot \exp(\pm y)$. The argument of the running coupling constant is taken as $\Omega^2 = \max(\kappa_1^2, \kappa_2^2, p_t^2)$.

Here we shall not discuss the distributions of "produced" gluons, which can be found in [ 11]. Instead we shall discuss what are typical values of $x_1$ and $x_2$ in the jet (particle) production. Average value $< x_1 >$ and $< x_2 >$, shown in Fig.1, only weakly depend on the model of UGD. For $y \sim 0$ at the RHIC energy W = 200 GeV one tests UGD's at $x_g = 10^{-3}$ - $10^{-2}$. When $|y|$ grows one tests more and more asymmetric (in $x_1$ and $x_2$) configurations. For large $|y|$ either $x_1$ is extremely small ($x_1 < 10^{-4}$) and $x_2 \to 1$ or $x_1 \to 1$ and $x_2$ is extremely small ($x_2 < 10^{-4}$). These are regions of gluon momentum fraction where the UGD's is rather poorly known. The approximation used in obtaining UGD's are valid certainly only for $x < 0.1$. [1] In order to extrapolate the gluon distribution to $x_g \to 1$ we multiply the gluon distributions from the previous section by a factor $(1-x_g)^n$, where n = 5-7.

## 3. From gluon to particle distributions

In Ref.[ 3] it was assumed, based on the concept of local parton-hadron duality, that the rapidity distribution of particles is identical to the rapidity distribution of

[1] an interesting discussion concerning the kinematics of the standard 2→2 and the 2→1 mechanism discussed here can be found in [ 18]




**Fig. 1.** $< x_1 >$ and $< x_2 >$ for $p_t > 0.5$ GeV and at W = 200 GeV.

gluons. In the present approach we follow a different approach which makes use of phenomenological fragmentation functions (FF's). For our present exploratory study it seems sufficient to assume $\theta_h = \theta_g$. This is equivalent to $\eta_h = \eta_g = y_g$, where $\eta_h$ and $\eta_g$ are hadron and gluon pseudorapitity, respectively. Then

$$y_g = \text{arsinh}\left(\frac{m_{t,h}}{p_{t,h}} \sinh y_h\right) \ , \tag{2}$$

where the transverse mass $m_{t,h} = \sqrt{m_h^2 + p_{t,h}^2}$. In order to introduce phenomenological FF's one has to define a new kinematical variable. In accord with $e^+e^-$ and $ep$ collisions we define a quantity $z$ by the equation $E_h = zE_g$. This leads to the relation

$$p_{t,g} = \frac{p_{t,h}}{z} J(m_{t,h}, y_h) \ , \tag{3}$$

where $J(m_{t,h}, y_h)$ is given in Ref.[ 11]. Now we can write the single particle distribution in terms of the gluon distribution as follows

$$\frac{d\sigma(\eta_h, p_{t,h})}{d\eta_h d^2 p_{t,h}} = \int dy_g d^2 p_{t,g} \int dz \ D_{g\to h}(z, \mu_D^2) \tag{4}$$
$$\delta(y_g - \eta_h) \ \delta^2\left(\vec{p}_{t,h} - \frac{z\vec{p}_{t,g}}{J}\right) \cdot \frac{d\sigma(y_g, p_{t,g})}{dy_g d^2 p_{t,g}} \ .$$

In the present calculation we use only LO FF's from [ 12] or [ 13].




**Fig. 2.** Charged-pion pseudrapidity distribution at W = 200 GeV for different models of UGD's. In this calculation $p_{t,h} > 0.2$ GeV. The experimental data of the UA5 collaboration are taken from [ 14].

Let us analyze now how the results for pseudorapidity distributions depend on the choice of the UGD. In Fig.2 I compare pseudorapidity distribution of charged pions for different models of UGD's. In this calculation FF from [ 12] has been used.

In contrast to Ref.[ 3], where the whole pseudorapidity distribution, including fragmentation regions, has been well described in an approach similar to the one presented here, in the present approach pions produced from the fragmentation of gluons in the $gg \to g$ mechanism populate only midrapidity region, leaving room for other mechanisms in the fragmentation regions. These mechanisms involve quark/antiquark degrees of freedom or leading protons among others. This strongly suggests that the agreement of the result of the $gg \to g$ approach with the PHOBOS distributions [ 2] in Ref.[ 3] in the true fragmentation region is rather due to approximations made in [ 3] than due to correctness of the reaction mechanism. In principle, this can be verified experimentally at RHIC by measuring the $\pi^+/\pi^-$ ratio in proton-proton scattering as a function of (pseudo)rapidity in possibly broad range. The BRAHMS experiment can do it even with the existing apparatus.

In Fig.3 we compare the theoretical transverse momentum distributions of charged pions obtained with different gluon distributions with the UA1 collaboration data [ 15]. The best agreement is obtained with the Karzeev-Levin gluon distribution. The distribution with the GBW model is much too steep in compar-

ison to experimental data. This is probably due to neglecting QCD evolution in [ 7].




**Fig. 3.** Transverse momentum distributions of charged pions at W = 200 GeV for BKK1995 FF and different UGD's. The experimental data are taken from [ 15].

## 4. Unintegrated parton distributions from the solution of the Kwieciński CCFM equations

Many unintegrated gluon distributions in the literature are ad hoc parametrizations of different sets of experimental data rather than derived from QCD. Recently Jan Kwieciński and collaborators [ 16, 17] have shown how to solve so-called CCFM equations by introducing unintegrated parton distributions in the space conjugated to the transverse momenta [ 16]

$$f_q(x,\kappa^2,\mu^2) = \frac{1}{2\pi}\int \exp\left(i\vec{\kappa}\vec{b}\right)\ \tilde{f}_q(x,b,\mu^2)\ d^2b\ . \tag{5}$$

The approach proposed is very convenient to introduce the nonperturbative effects like internal (nonperturbative) transverse momentum distributions of partons in nucleons. It seems reasonable to include the nonperturbative effects in the factorizable way

$$\tilde{f}_q(x,b,\mu^2) = \tilde{f}_q^{CCFM}(x,b,\mu^2)\cdot F_q^{np}(b)\ . \tag{6}$$




**Fig. 4.** Leading-order diagrams for parton production

In the following, for simplicity, we use a flavour and $x$-independent form factor

$$F_q^{np}(b) = F^{np}(b) = \exp\left(\frac{b^2}{4b_0^2}\right) \tag{7}$$

which describes the nonperturbative effects.

## 5. From unintegrated parton distributions to meson production

The $gg \to g$ mechanism considered in the literature is not the only one possible. In Fig.4 we show two other leading order diagrams. They are potentially important in the so-called fragmentation region. The formulae for inclusive quark/antiquark distributions are similar to formula (1) and will be given explicitly elsewhere [ 19]. The inclusive distributions of hadrons (pions, kaons, etc.) are obtained through a convolution of inclusive distributions of partons and flavour-dependent fragmentation functions

$$\frac{d\sigma(\eta_h, p_{t,h})}{d\eta_h d^2p_{t,h}} = \int_{z_{min}}^{z_{max}} dz \frac{J^2}{z^2}$$

$$D_{g\to h}(z,\mu_D^2) \frac{d\sigma^A_{gg\to g}(y_g, p_{t,g})}{dy_g d^2p_{t,g}}\Bigg|_{\substack{y_g=\eta_h \\ p_{t,g}=Jp_{t,h}/z}}$$

$$\sum_{f=-3}^{3} D_{q_f\to h}(z,\mu_D^2) \frac{d\sigma^{B_1}_{q_f g\to q_f}(y_{q_f}, p_{t,q_f})}{dy_{q_f} d^2p_{t,q}}\Bigg|_{\substack{y_q=\eta_h \\ p_{t,q}=Jp_{t,h}/z}}$$




**Fig. 5.** Pseudorapidity distribution of charged pions at W = 200 GeV calculated with the CCFM parton distributions. The experimental data are taken from [ 15]. The thin solid line is the gluon-gluon contribution while the dashed lines represent the gluon-(anti)quark and (anti)quark-gluon contributions. In this calculation gluon, (anti)quark fragmentation functions from [ 13] have been used.

$$\sum_{f=-3}^{3} D_{q_f\to h}(z,\mu_D^2)\frac{d\sigma^{B_2}_{gq_f\to q_f}(y_{q_f},p_{t,q_f})}{dy_{q_f}d^2p_{t,q}}\Bigg|_{\substack{y_q=\eta_h\\ p_{t,q}=Jp_{t,h}/z}}.$$

In Fig.5 we show the distribution in pseudorapidity of charged pions calculated with the help of the CCFM parton distributions [ 17] and the Gaussian form factor (7) with $b_0 = 0.5$ GeV$^{-1}$, adjusted to roughly describe the UA5 collaboration data. Now both gluon-gluon and (anti)quark-gluon and gluon-(anti)quark fussion processes can be included in one consistent framework. As anticipated the missing up to now terms are more important in the fragmentation region, although its contribution in the central rapidity region is not negligible. More details concerning the calculation will be presented elsewhere [ 19].

For completeness in Fig.6 we show transverse momentum distribution of positive and negative pions for different incident energies. The presence of diagrams $B_1$ and $B_2$ leads to an asymmetry in $\pi^+$ and $\pi^-$ production. The higher the incident energy the smaller the asymmetry. This is caused by the dominance of diagram $A$ at high energies.




**Fig. 6.** Transverse momentum distribution of $\pi^+$ (solid) and $\pi^-$ (dashed) mesons in proton-proton collisions for different center-of-mass energies.

## 6. Conclusions

We have calculated the inclusive distributions of gluons and associated charged $\pi$'s in the NN collisions through the $gg \to g$ mechanism in the $k_t$-factorization approach. The results for several UGD's proposed recently have been compared. The results, especially $p_{t,h}$ distributions, obtained with different models of UGD's differ considerably.

Contrary to a recent claim in Ref.[ 3], we find that the gluonic mechanism discussed in the literature does not describe the inclusive spectra of charged particles in the fragmentation region, i.e. in the region of large (pseudo)rapidities for any UGD from the literature. The existing UGD's lead to the contributions which almost exhaust the strength at midrapidities but leave room for other mechanisms in the fragmentation regions. We conclude that at presently available energies the gluonic mechanism is not the only contribution.

We propose new mechanisms, neglected so far in the literature, which involve also quark/antiquark degrees of freedom and are based on (anti)quark-gluon and gluon-(anti)quark fusion processes followed by the subsequent fragmentation. These missing mechanisms have been estimated in the approach based on unintagrated parton (gluon, quark, antiquark) distributions originating from the solution of a set of couple CCFM equations proposed recently by Kwieciński and coworkers. They lead to an asymmetry in the production of the $\pi^+$ and $\pi^-$ mesons. While at

lower energies of the order of 20 – 50 GeV such an asymmetry has been observed experimentally, it was not yet studied at RHIC. The BRAHMS collaboration at RHIC has a potential to study such asymmetries.

## References


1. Proceedings of the Quark Matter 2002 conference, July 2002, Nantes, France; Proceedings of the Quark Matter 2004 conference, January 2004, Oakland, USA.
2. B.B. Back et al.(PHOBOS collaboration), Phys. Rev. Lett. **87** (2001) 102303-1.
3. D. Kharzeev and E. Levin, Phys. Lett. **B523** (2001) 79.
4. I.G. Bearden et al.(BRAHMS collaboration), Phys. Rev. Lett. **87** (2001) 112305.
5. A.J. Askew, J. Kwieciński, A.D. Martin and P.J. Sutton, Phys. Rev. **D49** (1994) 4402.
6. K. Golec-Biernat and M. Wüsthoff, Phys. Rev. **D59** (1999) 014017.
7. K. Golec-Biernat and M. Wüsthoff, Phys. Rev. **D60** (1999) 114023-1.
8. M.A. Kimber, A.D. Martin and M.G. Ryskin, Eur. Phys. J. **C12** 655; M.A. Kimber, A.D. Martin and M.G. Ryskin, Phys. Rev. **D63** (2001) 114027-1.
9. J. Blümlein, a talk at the workshop on Deep Inelatic Scattering and QCD, hep-ph/9506403.
10. L.V. Gribov, E.M. Levin and M. G. Ryskin, Phys. Lett. **B100** (1981) 173.
11. A. Szczurek, Acta Phys. Polon. **34** (2003) 3191.
12. J. Binnewies, B.A. Kniehl, G. Kramer, Phys. Rev. **D52** (1995) 4947.
13. S. Kretzer, Phys. Rev. **D62** (2000) 054001.
14. G.J. Alner et al. (UA5 collaboration), Z. Phys. **C33** (1986) 1.
15. C. Albajar et al. (UA1 collaboration), Nucl. Phys. **B335** (1990) 261.
16. J. Kwieciński, Acta Phys. Polon. **B33** (2002) 1809.
17. A. Gawron, J. Kwieciński and W. Broniowski, Phys. Rev. **D68** (2003) 054001.
18. A. Accardi, these proceedings (nucl-th/0405046).
19. A. Szczurek and M. Czech, a paper in preparation.